

NICU NURSING CARE PLAN <29 weeks and/or <1000g FOR FIRST 72 Hours



Name:		DOB:			General Care/Measurements/Assessments			
		GA:		Check Q 4 - 6	hrs	We	ight Q 24 hrs.	, always with CPAP on
Birth Weight:		CGA:	Date:	Vitals Q 1hr			Length	NO
Mom (cell):		(home):		BP cuff Q check	ks, ONLY if no	UAL	HC	NO
Dad (cell):		(home):		Use only prem (white	te, non-adhesive) :	sat probe	Girth:	NO
NeoConnect:	Y/N	Siblings:		Change sat probe p	lacement Q full ch	neck	Other:	
Language: F □ E □ Other				Bath				
Diagnosis:				NO bath until 72hrs of life NO CHG wipe down				
				Tests/Consults		Dressings/Ostomies		
				MRSA/VRE/KPO	C: only if outbo	rn and > 24 hr	s of life	0
				NB screening blo	•			
Treatments/Operations:				NB screening urine due://				
Treatments, operations.				Other tests: Consults:				
				o iner tests.		Consums		
						All non-nurs	ing/non-family	Use neonatal bags (small
				Record all blood o			before accessing	preemie pouches, Hollister
Precautions:				Blood transfusion	•	be	aby	#3777 and 3778) only
				Temperature/Humidity/Skin Care				
Respiratory/Cardiovascular Ventilator/CPAP Parameters:				Humidity: <25 weeks - 85% ≥25 weeks - 75%				
		AD DDN M	L_	1		_	weeks - /5%	
Suctioning:	Min Q4h if on BCP		KO	Temp probe:	Tape with Me	•	. 1 . 1	
Oxygen:	Only one prong out				Change positi			1. 4. 1. / 11.
iNO:	Chest Physio:		l massage: NO	UAL/UVL:	Mepitac unde			directly on skin/rubbing
Saturation Targets: OWL. Allow desaturations 15-30 secs before \(\gamma \) FiO ₂				Developmental Care				
Elimination				To decrease stress:				
Strict input/output? Y Balance: Q 4 - 6hrs (with checks)				DOME CLOSED unless emergency.				
Foley → Inserted: Removed:				Use "Boost air curtain" (will last 20 mins) when accessing baby				
Balloon inflated: mL				• Cluster care				
Inform MD if urine output <0.5 ml/kg/hr or				See positioning principles at the back of care plan				
If urine output > hourly IV rate x 4hrs				Promote non-nutritive sucking				
Feeding				Keep dark (cover eyes during check)				
Type: PHM / Maternal breastmilk Calories: NO				Limit people/noise in room				
Route: OG				Use "ELG	GA sign-in shee	t" for all non-	nursing/non-fa	mily handling of baby
Volume:	Frequency:	Incre	ease:	• Teach pare	ents how to touc	ch		
Over:h	Flush	with 0.5ml sterile	water ONLY	To decrease risk	c of IVH:			
Feeding notes: Start trophic feeds within 12 hours of life				• Keep head midline x first 72 hours = NO Prone/Lateral				
	OIT Q check			• HOB at 30	0°			
OG→ Inserted:		Secured @		 Avoid rapi 	id position chan	iges		
Change due:						Teaching wit	h Parents	
Replogle → Inserte	ed: S	Secured @	_ To LWS □	☐ Hand hygien	ne 🗆	Manual expre	ession	□ Touch
	Fluid	ls		□ Books for ba	abies 🗆	Breast pump	use/cleaning	
TFI:	Weight:	Tota	l ml/day:	□ Preemie time	eline 🗆	Modified Ka	ngaroo Care	
	Lines							
				UAL Sutured	1@	UV	L Sutured	@ Single/
				Inserted	l:		Central/F	Peripheral Double
				Remove	ed:		Inserted:	
							Removed	l:
				Cleanse skin after insertion with sterile NS to remove any leaked CHG				
	PICC Externa	al Portion:		Central/F	Peripheral Single/			
	Inserted	l:		Removed	- 0			
				Dressin	g due:			
				Special Notes				
						-		
Updated by:		Date:		1				

<29 weeks and/or <1000g FOR FIRST 72 hours



Alignment

Flexion

Symmetry

Movement

Containment



SUPINE

- Weight appropriate containment device, legs inside (snuggle up size XS)
- Appropriate size rolls to provide 360° containment (stockinette roll)
- Appropriate size roll under knees to provide support (dry wipes)
- * No bundling









