



## Indications

- The Comfort Zone is a bundle of four strategies that have been shown to decrease pain and distress associated with needle procedures when used together:
  - Comfort positioning (kangaroo care, bundling for those up to 6 months of age, or facilitated tucking in the premature infant that cannot be bundled)
  - Direct breastfeeding or sucrose administration (\*\*Note: breastmilk alone, without the skin to skin contact, does NOT act as a replacement for sucrose)
  - Distraction (Eg. pacifier, music, talking to baby gently, patting or stroking baby)
  - Topical anesthetic (Lidocaine 4% cream)

## **Key points**

- All needle procedures are included, but indications for each strategy may differ depending on procedure (see table below).
- Contraindications for lidocaine 4% include allergy to lidocaine and urgent procedure (eg. Stat IV)

Procedure	Recommended Pain Management
Heel Prick	<ul> <li>Breastfeeding OR pacifier + sucrose 24%</li> <li>Comfort positioning</li> <li>Distraction</li> </ul>
Injection (SC or IM)	<ul> <li>Topical anesthetic (lidocaine 4%), covered with Tegaderm, 20-30 minutes prior to procedure start (refer to lexicomp or Lidocaine protocol for dosing)</li> <li>Breastfeeding OR pacifier + sucrose 24%</li> <li>Comfort positioning</li> <li>Distraction</li> </ul>
Lumbar Puncture	<ul> <li>Topical anesthetic (lidocaine 4%), covered with Tegaderm, 20-30 minutes prior to procedure start (refer to Lexicomp or Lidocaine protocol for dosing)</li> <li>Breastfeeding OR pacifier + sucrose 24%</li> <li>Distraction</li> </ul>
Venipuncture/ IV insertion	<ul> <li>Topical anesthetic (lidocaine 4%), covered with Tegaderm, 20-30 minutes prior to procedure start (refer to Lexicomp or Lidocaine protocol for dosing)</li> <li>Breastfeeding OR pacifier + sucrose 24%</li> <li>Comfort positioning (swaddling)</li> <li>Distraction</li> </ul>

## **DECEMBER 2019**

Page | 1 of 1