

FEEDING CRITERIA FOR BANKED BREAST MILK BASED ON HÉMA-QUÉBEC INVENTORY LEVELS

CRITERIA	- RED -	- ORANGE -	- YELLOW -	- GREEN -
CRITERIA INCLUSION: AGE AND WEIGHT	Less than 29 weeks (28 6/7 and less) OR < 1000g	Less than 30 weeks (29 6/7 and under) OR < 1300g	Less than 31 weeks (30 6/7 and less) OR < 1500g	Less than 32 weeks (31 6/7 and less) OR < 1750g
DURATION : AGE AND WEIGHT	Up to: 32 weeks OR 2000g	Up to: 32 weeks OR 2000g	Up to: 33 weeks corrected age OR 2000g OR Stop before according to treatment team	Up to: 34 weeks corrected age OR 2000g OR Stop before according to treatment team
NEC	- Post-NEC refeeding stage 2 or more - For 2 weeks once FPO	- Post-NEC refeeding stage 2 or more - For 2 weeks once FPO	- Post-NEC refeeding or Severe enteritis (grade 2 or higher) - For 2 weeks once FPO	- Post-NEC refeeding or Severe enteritis (grade 2 or higher) - For 3 weeks once FPO
POST-SURGICAL INTESTINAL PATHOLOGY	N/A	N/A	- Post omphalocele, gastroschisis, volvulus, diaphragmatic hernia, short gut - For 2 weeks once FPO	- Post omphalocele, gastroschisis, volvulus, diaphragmatic hernia, shortgut - For 3 weeks once FPO
MEDICAL INTESTINAL PATHOLOGY AND OTHER PATHOLOGIES	N/A	N/A	- Significant Cardiac Malformation - For 2 weeks once FPO	- Significant Cardiac Malformation - For 3 weeks once FPO - During cooling, stop once rewarmed.

BANKED BREAST MILK IS GIVEN TO REPLACE OR SUPPLEMENT THE CHILD'S MOTHER'S BREAST MILK